

IUSO Clinical Abbreviation List

| | |
|----------|---------------------------------------------------|
| Δ | Prism |
| Δ | Change |
| @ | at |
| 1° | Primary |
| 2° | Secondary |
| 3° | Tertiary |
| A | Assessment |
| A/V | Arteriole/Venule Ratio |
| A/C | Anterior Chamber |
| ACE | Angiotensin Converting Enzyme |
| ACIOL | Anterior Chamber Intraocular Lens |
| Adapt | Adaptation Discussed |
| ADHD | Attention Deficit Hyperactive Disorder |
| AION | Anterior Ischemic Optic Neuropathy |
| All | Allergy |
| ALT | Alternating |
| Argon LT | Argon Laser Trabeculoplasty |
| AM | Morning |
| AMD | Age Related Macular Degeneration |
| AMPPE | Acute Multifocal Posterior Pigment Epitheliopathy |
| ANA | Antinuclear Antibody |
| Ant | Anterior |
| APD | Afferent Pupillary Defect |
| ARMD | Age Related Macular Degeneration |
| ASAP | As Soon As Possible |
| ASTIG | Astigmatism |
| ATs | Artificial tears |
| BD | Base Down |
| BDR | Background Diabetic Retinopathy |
| BF | Bifocal |
| BI | Base In |
| bid | 2X a day |
| BL | Baseline |
| BO | Base Out |
| BRAO | Branch Retinal Artery Occlusion |
| BRVO | Branch Retinal Vein Occlusion |
| BS | Blood Sugar |
| BU | Base Up |
| BV | Binocular Vision |
| BXC | Binocular Crossed Cylinder |
| c | With |
| C/D | Cup to Disk Ratio |
| C/F | Cell & Flare |
| C/S | Conjunctiva/Sclera |
| C1 | Cyclogyl 1% |
| CA | Cancer, Carcinoma |
| CAI | Carbonic Anhydrase Inhibitor |
| CAT | Cataract |
| CB | Ciliary Body |
| cc | With Correction |
| CC | Chief Complaint |
| CF | Count Fingers |
| CF&F | Central Fix And Follow |
| CHF | Congestive Heart Failure |

| | |
|---------|---------------------------------------|
| CL | Contact Lens |
| CLARE | Contact Lens Associated Red Eye |
| CLPU | Contact Lens Peripheral Ulcer |
| CLR | Clear |
| cm | Centimeters |
| CME | Cystoid Macular Edema |
| CNVM | Choroidal Neovascular Membrane |
| c/o | Complaint of |
| COAG | Chronic Open Angle Glaucoma |
| CONF | Confrontation |
| Conj | Conjunctiva |
| COPD | Chronic Obstructive Pulmonary Disease |
| CPM | Continue Present Meds |
| CRAO | Central Retinal Artery Occlusion |
| CRVO | Central Retinal Vein Occlusion |
| CSME | Clinically Significant Macular Edema |
| CT | Cover Test |
| CT Scan | Computerized Tomography Scan |
| CVA | Cerebrovascular Accident |
| CVF | Confrontation Visual Field |
| CW | Continuous Wear |
| CWS | Cotton Wool Spot |
| CYL | Cylinder |
| d | Day |
| D | Disk, Diopter |
| D/W | Discussed With |
| D/B | Dot & Blot Hems |
| D/C | Discontinue |
| DD | Disk Diameter |
| DDx | Differential Diagnosis |
| DEE | Dilated Eye Exam |
| DES | Dry Eye Syndrome |
| Dev | Development |
| DFE | Dilated Fundus Exam |
| D/I | Dilation and Irrigation |
| DM | Diabetes Mellitus |
| DMP | Diabetes Mellitus Program |
| DNA | Did Not Answer |
| DNKA | Did Not Keep Appointment |
| Doxy | Doxycycline |
| DR | Diabetic Retinopathy |
| DS | Diopters Sphere |
| DTR | Deep Tendon Reflex |
| DVA | Distance Visual Acuity |
| DVMP | Disc, Vessels, Macula, Periphery |
| DW | Daily Wear |
| Dx | Diagnosis |
| Dz | Disease |
| ECCE | Extracapsular Cataract Extraction |
| ECP | Endo Cyclo Photocoagulation |
| Ed | Education |
| EKC | Epidemic Keratoconjunctivitis |
| ELISA | Enzyme-Linked Immunosorbent Assay |

IUSO Clinical Abbreviation List

| | |
|---------|-------------------------------------------------|
| Endo | Endothelium |
| EOM | Extraocular Muscles/ Motility |
| EOM/I | Extraocular Motility Intact |
| EP | Esophoria |
| EP' | Esophoria @ Near |
| Epi | Epithelium |
| ER | Emergency Room/ Department |
| ERM | Epi-retinal Membrane |
| ESR | Erythrocyte Sedimentation Rate |
| ET | Esotropia |
| ET' | Esotropia @ near |
| ETOH | Ethanol/ Alcohol |
| Eval | Evaluation |
| EW | Extended Wear |
| EXT | External |
| F | Father |
| F & F | Fix and Follow |
| F/U | Follow-Up |
| FA | Fluorescein Angiogram |
| FANG | Fluorescein Angiogram |
| FAZ | Foveal Avascular Zone |
| FB | Foreign Body |
| FBS | Fasting Blood Sugar |
| FHx/ FH | Family History |
| FLR | Foveal Light Reflex |
| FTA-ABS | Fluorescent Treponemal Antibody Absorption Test |
| FTFC | Full To Count Fingers |
| GC | Gonococcus |
| GCA | Giant Cell Arteritis |
| GF | Grandfather |
| Glauc | Glaucoma |
| GLC | Glaucoma |
| GM | Grandmother |
| GP | Gas Permeable |
| GPC | Giant Papillary Conjunctivitis |
| gtt | Drop(s) |
| GTT | Glucose Tolerance Test |
| h | Hour |
| H/B/T | Holes/Breaks/Tears |
| h/o | History of |
| HA | Headache |
| HbA1C | Hemoglobin A1C |
| HLA | Human Leucocytic Antigen |
| HM | Hand Motion |
| HSK | Herpes Simplex Keratitis |
| HSV | Herpes Simplex Virus |
| HT | Hypertropia |
| HTN | Hypertension |
| HVF | Humphrey's Visual Field Analyzer |
| Hx | History |
| HZV | Herpes Zoster Virus |
| I/L | Iris/Lens |
| ICCE | Intracapsular Cataract Extraction |
| ICE | Iridocorneal Epithelial Syndrome |

| | |
|-------|------------------------------------------------------------------------------|
| IGT | Impaired Glucose Tolerance |
| IK | Interstitial Keratitis |
| INO | Internuclear Ophthalmoplegia |
| INT | Internal |
| IO | Inferior Oblique |
| IOL | Intraocular Lens |
| ION | Ischemic Optic Neuropathy |
| IOP | Intraocular Pressure |
| IRMA | Intraretinal Microvascular Abnormality |
| IRT | In Regards To |
| ISNT | Inferior Superior Nasal Temporal (rule used to assess optic disk appearance) |
| K | Cornea |
| KC | Keratoconjunctivitis |
| KCN | Keratoconus |
| KP | Keratic Precipitates |
| L | Left |
| LASIK | Laser InSitu Keratomileusis |
| L/L | Lids/Lashes |
| LKP | Lamellar Keratoplasty |
| L LL | Left Lower Lid |
| LP | Light Perception |
| LPI | Laser Peripheral Iridotomy |
| LR | Lateral Rectus |
| LUL | Left Upper Lid |
| LWOBS | Left Without Being Seen |
| M | Male |
| (M) | Maternal, Mother |
| M1 | Mydracyl 1% |
| MA | Microaneurysm |
| MCAR | Mires Clear & Regular |
| MEDS | Medications |
| MEWDS | Multiple Evanescent White Dot Syndrome |
| MG | Myasthenia Gravis |
| MGD | Meibomian gland disease/dysfunction |
| MGF | Maternal Grandfather |
| MGM | Maternal Grandmother |
| MK | Microbial Keratitis |
| mm | Millimeter |
| mmHg | Millimeters of Mercury |
| Mo(s) | Month(s) |
| MR | Medial Rectus |
| MRI | Magnetic Resonance Imaging |
| MRx | Manifest Refraction |
| MS | Multiple Sclerosis |
| MVA | Motor Vehicle Accident |
| N | Nasal |
| N/A | Not Available; Not applicable |
| NaFl | Sodium Fluorescein |
| NAG | Narrow Angle Glaucoma |
| NAP | No Apparent Pathology |
| NCT | Non-Contact Tonometry |
| NFL | Nerve Fiber Layer |

IUSO Clinical Abbreviation List

| | |
|--------|--------------------------------------------------------------|
| NI | Not Indicated |
| NIPH | No Improvement with Pinhole |
| NKDA | No Known Drug Allergy |
| NKMA | No Known Medical Allergy |
| NL | Normal |
| NLDO | Naso-lacrimal Duct Obstruction |
| NLP | No Light Perception |
| NPA | Near Point of Accommodation |
| NPATs | Nonpreserved Artificial Tears |
| NPC | Near Point of Convergence |
| NPDR | Non-Proliferative Diabetic Retinopathy |
| NS | Nuclear Sclerosis, |
| NSAID | Non-Steroidal Anti-Inflammatory Drug |
| NTG | Normal Tension Glaucoma |
| NVA | Near Visual Acuity |
| NVD | Neovascularization at the Disk |
| NVE | Neovascularization Elsewhere |
| NVG | Neovascular Glaucoma |
| NVI | Neovascularization of the Iris |
| NVO | Near Vision Only |
| O | Objective |
| O | Single Surround HOTV |
| ⊕ | Orthophoria |
| OAG | Open Angle Glaucoma |
| OCT | Ocular Coherence Tomography |
| OD | Right Eye |
| OKN | OptoKinetic Nystagmus |
| ON | Optic Nerve |
| ONH | Optic Nerve Head |
| OR | Over Refraction |
| OS | Left Eye |
| OU | Both Eyes |
| P | Plan |
| (P) | Paternal |
| Pach | Pachymetry |
| PAL | Progressive Addition Lens |
| PAS | Peripheral Anterior Synechia |
| pc | After Meals |
| PC | Primary Care |
| PCF | PharyngoConjunctival Fever |
| PCIOL | Posterior Chamber Intraocular Lens |
| PCM | Primary Care Manager |
| PCO | Posterior Capsular Opacification |
| PCP | Primary Care Provider |
| PD | Pupillary Distance |
| PDR | Proliferative Diabetic Retinopathy |
| PE2.5 | Phenylephrine 2.5% |
| PERIPH | Periphery |
| PERRLA | Pupils Equally Round and Reactive to Light and Accommodation |
| PHACO | Phacoemulsification |
| PH | Pinhole |
| PI | Peripheral Iridectomy/ Iridotomy |
| PKC | Phlyctenular Keratoconjunctivitis |

| | |
|----------|----------------------------------------------------|
| PLO | Plano |
| PM | Afternoon/Night (12noon – 12 midnight) |
| PMD | Pellucid Marginal Degeneration |
| PMMA | PolyMethylMethAcrylate |
| po | By Mouth |
| POAG | Primary Open Angle Glaucoma |
| POHS | Presumed Ocular Histoplasmosis Syndrome |
| POLYCARB | Polycarbonate Lens |
| Post | Posterior |
| PR | Poor Patient Responses |
| prev | Previous |
| PRH | Preretinal Hemorrhage |
| PRK | Photorefractive Keratectomy |
| prn | As Needed |
| prog | Prognosis |
| PRP | Pan-Retinal Photocoagulation |
| PSC | Posterior Subcapsular Cataract |
| Pt | Patient |
| PVD | Posterior Vitreous Detachment |
| PXE | PsuedoXanthoma Elasticum |
| PXF | PsuedoExfoliation |
| q | each, every |
| Q&D | Quiet and Deep |
| q2h | Every Two Hours |
| qh | Every Hour |
| qhs | At Bedtime |
| qid | Four Times a Day |
| qod | Every Other Day |
| R | Right |
| Ret | Retinoscopy |
| R & R | Recess & Resect |
| R/B//A | Risks, Benefits, Indications, Alternatives |
| R/O | Rule Out |
| RA | Rheumatoid Arthritis |
| RBS | Random Blood Sugar |
| RD | Retinal Detachment |
| Ref | Refraction |
| RFV | Reason for Visit |
| RGP | Rigid Gas Permeable Contact Lens |
| RK | Radial Keratotomy |
| RLL | Right Lower Lid |
| ROP | Retinopathy Of Prematurity |
| RP | Retinitis Pigmentosa |
| RPE | Retinal Pigment Epithelium |
| RSVP | Redness, Sensitivity to Light, Vision Change, Pain |
| RTC | Return To Clinic |
| RUL | Right Upper Lid |
| Rx | Refraction/ Prescription/ Therapy |
| rxn | Reaction |
| S | Subjective/ Signs |
| S | Single Surround Snellen |
| (SB) | Sibling |

IUSO Clinical Abbreviation List

| | |
|-------|--------------------------------------------|
| s/p | Status Post |
| s/s | Signs and Symptoms |
| sc | Without Correction |
| SCL | Soft Contact Lens |
| SEI | Sub-epithelial Infiltrate |
| SHx | Social History |
| sig | Label |
| SITA | Swedish Interactive Thresholding Algorithm |
| SL | Schwalbe's Line |
| SLACH | Soft Lens Associated Corneal Hypoxia |
| SLEx | Slit Lamp Exam |
| SLK | Superior Limbic Keratoconjunctivitis |
| SLT | Selective Laser Trabeculoplasty |
| SO | Superior Oblique |
| SPH | Sphere |
| SPK | Superficial Punctate Keratitis |
| SR | Superior Rectus |
| SRNVM | Sub-Retinal Neovascular Membrane |
| SRx | Spectacle Prescription or Glasses |
| SS | Scleral Spur |
| STAT | Immediately |
| SVP | Spontaneous Venous Pulsation |
| SWAP | Short Wavelength Automated Perimetry |
| Sx | Symptoms |
| Sz | Seizure |
| T | Tight (used to describe contact lens fit) |
| T1 | Tropicamide 1% |
| TA | Applanation Tonometry |
| TBUT | Tear Break-Up Time |
| TCN | Tetracycline |
| TDO | Tear Duct Obstruction |
| TF | Trial Frame |
| TIA | Transient Ischemic Attack |
| tid | Three Times a Day |
| TM | Trabecular Meshwork |
| Tp | Tonopen |
| Tx | Treatment |
| ung | Ointment |
| V | Vessels |
| VA | Visual Acuity |
| VDRL | Venereal Disease Research Laboratory Test |
| VEP | Visually Evoked Potential |
| VF | Visual Field |
| VH | Vitreous Hemorrhage |
| VI | Visual Impairment |
| Vit | Vitreous |
| VT | Visual Therapy |
| w/ | With |
| w/o | Without |
| WCP | warm compress |
| WI | Walk-In |

| | |
|-----|-------------------------------|
| Wk | Week |
| WNL | Within Normal Limits |
| X | by/for |
| XP | Exophoria |
| XP' | Exophoria @ Near |
| XT | Exotropia |
| XT' | Exotropia @ Near |
| YO | Year Old |
| YAG | Yttrium-Aluminum-Garnet Laser |
| Yr | Year |